

THE PARKES PULSAR TIMING ARRAY PROJECT

GEORGE HOBBS

CSIRO Astronomy and Space Science, PO Box 76, Epping, NSW, 1710, Australia

E-mail: george.hobbs@csiro.au

(Received November 30, 2014; Revised May 31, 2015; Accepted June 30, 2015)

ABSTRACT

The main goals of the Parkes Pulsar Timing Array (PPTA) project are to 1) detect ultra-low-frequency gravitational waves, 2) improve the solar system planetary ephemeris and 3) provide a long-term, stable time standard. In this paper, we highlight the main results from the project so far and discuss our expectations for the future.

Key words: pulsars: general

1. INTRODUCTION

Very precise determinations of pulse times-of-arrival (ToAs) from millisecond pulsars can be obtained with the Parkes radio telescope. The pulsar timing method (see e.g., Edwards et al., 2006) provides the means by which those ToAs can be modelled using a relatively simple parameterisation of the pulsar and the interstellar medium. Physical effects that are not in the parameterisation, but cause variations in the pulse ToAs, will lead to structure in the “timing residuals” (the difference between the observed and predicted ToAs). Phenomena that are local to the pulsar (e.g., magnetospheric variations) will induce timing residuals that are not correlated between different pulsars. Errors in the clock used for measuring the ToAs will lead to identical residuals for all pulsars analysed with that particular clock. The effect of errors in the solar system ephemeris or the passage of gravitational waves (GWs) past the Earth will lead to timing residuals that can be distinguished by studying how the timing residuals are correlated on different angular scales.

The Parkes Pulsar Timing Array (PPTA) was started in 2004 (see Manchester et al. 2013a for details and Hobbs 2012 for a historical review) and has the three main goals of 1) detecting GWs, 2) improving the solar system ephemerides and 3) developing a pulsar-based timescale. In this paper we describe our research related to these main project goals, but note that the data sets have also been used for numerous other projects such as studying the interstellar medium (You et al., 2007; Keith et al., 2013), the solar wind (You et al., 2012) and the pulsars themselves (Osłowski et al., 2011; Shannon et al., 2014; Verbiest et al., 2009).

2. THE PPTA DATA SETS

The PPTA project has been described in detail by Manchester et al. (2013a). In brief, all observations are obtained with the 64-m diameter Parkes radio telescope. We now observe 24 millisecond pulsars (some pulsars have been recently added, other pulsars have been removed from the sample) with a typical observing cadence of about three weeks. During each observing session we attempt to observe each pulsar at least once with a 20 cm receiver that provides a bandwidth of 256 MHz and once with a dual-frequency receiver that simultaneously enables observations in the 10 and 50 cm bands. In the 10 cm band, 1 GHz of bandwidth is available and the 50 cm band provides a bandwidth of 64 MHz. We currently use two digital filterbank systems (PDFB3 and PDFB4) to record the data along with two coherent dedispersion instruments (CASPSR and APSR). Typical observation times are 1 hour and each observation is preceded by a short calibration observation of a pulsed noise signal.

We process the observations using the PSRCHIVE software package (Hotan, van Straten & Manchester, 2004) to obtain pulse ToAs. Those ToAs are subsequently processed using the TEMPO2 software package (Hobbs et al., 2006).

High-quality PPTA observations started in the year 2005. Data from 2005 to 2011 have been published as the first data release (known as the “dr1” data set; see Section 7 and the timing residuals in Figure 2). Combining these data with earlier observations from the Parkes telescope (Verbiest et al. 2009) led to the extended data release 1 (known as “dr1e”). For some pulsars this provides a data span extending back to the year 1996.

3. NEW TECHNIQUES

In order to achieve the timing precision necessary to search for GWs, we have had to make numerous im-




Figure 1. The sensitivity of the PPTA (red circle), space-based detectors, such as eLISA, and ground-based detectors, such as LIGO, to gravitational waves. Pulsar timing arrays are sensitive to GWs with frequencies between $\sim 10^{-9}$ to 10^{-7} Hz.

improvements to our software and algorithms. van Straten (2013) describes calibration methods that are required to determine ToAs at this level of precision. Previous timing software packages did not provide the necessary level of precision and so the TEMPO2 software package (Hobbs et al., 2006) was developed with the aim of including all phenomena that affect pulse ToAs at the >1 ns level. Timing residuals for pulsars that have been monitored over many years are affected by low-frequency noise processes. Coles et al. (2011) described how to generalise the least-squares-fitting routines within TEMPO2 to account for such noise. Deng et al. (2012) discussed how pulsar timing data sets could be interpolated and extrapolated.

The main project goal requires us to search for GWs, errors in time standards and errors in the solar system ephemeris. To do this we have introduced into TEMPO2 the ability to fit for functions of arbitrary shapes that have a monopolar, dipolar or a quadrupolar spatial signature. These can be fitted globally to the entire data set simultaneously with a fit for the specific timing model parameters for each individual pulsar. This has required the development of a global (introduced in Champion et al., 2010) and constrained (Keith et al., 2013) least-squares fitting procedure.

The dominant signal seen in the timing residuals for most pulsars is caused by dispersion measure variations. A procedure to remove those variations without removing any underlying GW signal was described by Keith

et al. (2013).

Oslowski et al. (2011, 2013) and Shannon et al. (2014) showed that the timing residuals for some of the PPTA pulsars are affected by jitter noise. This jitter noise was thought to be a fundamental noise floor in pulsar timing. However, Oslowski et al. (2011) demonstrated a method that led to an improvement in the rms timing residuals for PSR J0437–4715 of ~ 20 per cent. An improved version of this method doubled this improvement to ~ 40 per cent.

4. GRAVITATIONAL WAVES

Detweiler (1979) showed that GWs with frequencies between 10^{-9} – 10^{-7} Hz induce timing residuals that could be detectable with a sufficiently sensitive PTA. As shown in Figure 1 these GW frequencies are complementary to those probed by other GW detectors such as LIGO/Virgo (ground-based) and eLISA (space-based). The astrophysical predictions shown on this figure were obtained at the start of the PPTA project. More recent research has been published by Sesana et al (2008; 2009) and Ravi et al. (2014b). They show that the most likely source of detectable GWs will arise from a large number of coalescing supermassive binary black hole systems at the centres of merging galaxies¹. Ravi et al. (2014b)

¹Sanidas et al. (2012) have also described implications of recent GW limits for models of cosmic strings. Some such models produce a GW background that has a higher amplitude to the predicted backgrounds from black hole coalescence.

predicts a “most likely” signal, which is indicated by the solid green line in Figure 1. The red circle indicates our most recent bound on the amplitude of the GW background (Shannon et al. 2013) showing that we have already ruled out many of the early predictions.

The design of the PPTA experiment (number of pulsars, observing cadence etc.) was based on Jenet et al. (2005) who showed that it is necessary to observe around 20 pulsars, with weekly observing cadence over a duration of ~ 5 yr to make a detection of the GW background. Even though we have searched for such a background (e.g., Yardley et al., 2011), no detection has yet been made. Instead we have continued to place ever more stringent constraints on the amplitude of any possible background (Jenet et al., 2006; Shannon et al., 2013). The most recent limits (Shannon et al., 2013) rule out some models for black-hole formation and evolution in galaxies.

We have also searched for individual supermassive binary black hole systems (Yardley et al., 2010 and Zhu et al., 2014). Our most recent work shows that, in some sky directions, we are sensitive to GWs from circular supermassive binary black holes with masses of $10^9 M_\odot$ out to a luminosity distance of about 100 Mpc. We have also searched for the burst event predicted from the final merger of an individual black hole binary system. Wang et al. (submitted to MNRAS) places bounds on the size of any such events and shows that the expected event rate for detectable bursts is so low that our non-detection is not surprising.

Our algorithms have been based on frequentist statistics. In the future we will continue to develop our algorithms and apply them to increasingly sensitive data sets. We will also be applying the Bayesian methods developed elsewhere (e.g., van Haasteren et al., 2011; Demorest et al., 2013; Lentati et al., 2014) to our data sets. Finally, the detection of a GW signal will be of enormous scientific interest. We are currently carrying out research to ensure that our algorithms will not give a false positive detection.

5. THE SOLAR SYSTEM EPHEMERIS

The pulsar timing method relies on knowledge of the observatory position with respect to the solar system barycentre. In Champion et al. (2010) we used observations from the PPTA combined with observations from the Arecibo and Effelsberg telescopes in order to search for errors in the assumed solar system barycentric position caused by inaccurate planetary mass estimates. This technique allowed us to provide very precise estimates of the planetary masses. These masses were shown to be consistent with independent space-craft determinations.

The Champion et al. (2010) method can find mass errors in known planetary objects, but cannot be applied to search for unknown objects in the solar system. We have updated the TEMPO2 software package to fit for errors in the Earth–solar system barycentric vector as a function of time. That method is currently being applied to the PPTA data in order to search for unknown




Figure 2. The timing residuals from the first PPTA data release. The value gives the range of the residuals for a given pulsar in microseconds. Figure from Wang et al. (submitted to MNRAS).

objects. The results of this processing will be published elsewhere.

Deng et al. (2013) assumed that the position of the solar system barycentre was known and developed a procedure to use observations of millisecond pulsars to determine the position of the telescope. Of course, the telescope need not be on Earth and we demonstrated that with an X-ray telescope on board a spacecraft on a trajectory between Earth and Mars it is possible to determine its position with a precision and accuracy of ~ 10 km .

6. A PULSAR-BASED TIME STANDARD

Hobbs et al. (2012) and references therein demonstrated how errors in a terrestrial time standard could be identified by searching for the common signal in the pulsar timing residuals. Any such errors could then be “corrected” using the pulsar data enabling a long-term, stable time scale using both atomic clocks and pulsar observations. We demonstrated that known changes to International Atomic Time (TAI) could be identified using the PPTA data set and there was evidence for possible discrepancies with the world’s best time standard, terrestrial time as realised by the BIPM.

7. DATA ACCESS

The raw observations and corresponding calibration data obtained for the project are available from the CSIRO data archive (data.csiro.au; Hobbs et al., 2011). All data are publically available for download after an embargo period of 18 months. Making use of these data requires knowledge of the processing software (such as PSRCHIVE) to calibrate and visualise the data.

We have also made pulse ToAs, timing parameters and red-noise models available. Our first data release of processed data is also available from the data archive². We are currently working towards a second data release, which we expect will include all observations until mid-2014 and should be published by the end of this year.

8. THE INTERNATIONAL PULSAR TIMING ARRAY AND FUTURE TELESCOPES

Over the next few years it is expected that the Parkes telescope will be upgraded. We are hoping that a new ultra-wide band receiver (covering from around 700 MHz to 3 GHz) will be commissioned along with a new coherent-dedispersion backend system to process and record the data.

The PPTA data set is already being combined with data from the European (Kramer & Champion, 2013) and North American (McLaughlin 2013) timing arrays to form the International Pulsar Timing Array (IPTA) data set (Manchester et al., 2013b). The IPTA data set will be more sensitive to GWs, ephemeris errors and clock errors than the individual PTA data sets and it is likely that, in the relatively near future, an IPTA data set will lead the first detection of GWs.

In the longer term new telescopes are being designed and built. The major new telescopes for PTA research will be the Five Hundred Meter Spherical Telescope (FAST) being built in China (see Hobbs et al., 2014 for a review of PTA research on FAST), MeerKAT in South Africa and the Square Kilometre Array (SKA; Lazio, 2013).

9. CONCLUSIONS

The PPTA project has made significant progress towards its three major goals. With future upgrades to the Parkes telescope, the PPTA data sets should remain the most sensitive PTA data sets in the Southern Hemisphere until observations begin with the SKA Phase 1. Combining these observations with similarly sensitive data sets from the Northern Hemisphere should lead to the first GW detection, whilst enabling incredibly varied research into topic such as the millisecond pulsar population, interstellar medium, solar corona, spacecraft navigation, clocks and planets.

ACKNOWLEDGMENTS

The Parkes radio telescope is part of the Australia Telescope, which is funded by the Commonwealth of Australia for operation as a National Facility managed by the Commonwealth Scientific and Industrial Research Organisation (CSIRO). GH acknowledges support from the Australian Research Council (ARC) Future Fellowship. GH thanks R. Manchester, M. Kerr, X. Zhu and S. Osłowski for useful comments on the manuscript.

REFERENCES

- Champion, D., Hobbs, G., & Manchester, R. N., et al., 2010, Measuring the Mass of Solar System Planets Using Pulsar Timing, *ApJ*, 720, 201
- Coles, W., Hobbs, G., & Champion, D. J., et al., 2011, Pulsar Timing Analysis in the Presence of Correlated Noise, *MNRAS*, 418, 561
- Detweiler, S., 1979, Pulsar Timing Measurements and the Search for Gravitational Wwaves, *ApJ*, 234, 1100
- Demorest, P. B., Ferdman, R. D., & Gonzalez, M. E., et al., 2013, Limits on the Stochastic Gravitational Wave Background from the North American Nanohertz Observatory for Gravitational Waves, *ApJ*, 762, 94
- Deng, X. P., Coles, W., & Hobbs, G. B. et al., 2012, Optimal Interpolation and Prediction in Pulsar Timing, *MNRAS*, 424, 244
- Deng, X. P., Hobbs, G. B., & You, X. P., 2013, Interplanetary Spacecraft Navigation Using Pulsars, *AdSpR*, 52, 1602
- Edwards, R. T., Hobbs, G. B., & Manchester, R. N., 2006, TEMPO2, a New Pulsar Timing Package - II. The Timing Model and Precision Estimates, *MNRAS*, 372, 1549
- Hobbs, G., Edwards, R., & Manchester, R. N., 2006, TEMPO2, a New Pulsar-timing Package - I. An Overview, *MNRAS*, 369, 655
- Hobbs, G., et al., 2011, The Parkes Observatory Pulsar Data Archive, *PASA*, 28, 202
- Hobbs, G., Coles, W., & Manchester, R. N., et al., 2012, Development of a Pulsar-based Time-scale, *MNRAS*, 427, 2780

²<http://dx.doi.org/10.4225/08/534CC21379C12>

- Hobbs, G., 2012, The Parkes Pulsar Timing Array: What We've Done and What We're Doing, arXiv, 1210, 0977
- Hobbs, G., Dai, S., & Manchester, R. N., et al., 2014, The Role of FAST in Pulsar Timing Arrays, arXiv, 1407, 0435
- Hotan, A., van Straten, W., & Manchester, R. N., 2004, PSRCHIVE and PSRFITS: An Open Approach to Radio Pulsar Data Storage and Analysis, PASA, 21, 302
- Jenet, F. A., Hobbs, G. B., Lee, K. J., & Manchester, R. N., 2005, Detecting the Stochastic Gravitational Wave Background Using Pulsar Timing, ApJ, 625, L123
- Jenet, F. A., Hobbs, G. B., & van Straten, W., et al., 2006, Upper Bounds on the Low-Frequency Stochastic Gravitational Wave Background from Pulsar Timing Observations: Current Limits and Future Prospects, ApJ, 653, 1571
- Keith, M., Coles, W., & Hobbs, G. B., et al., 2013, Measurement and Correction of Variations in Interstellar Dispersion in High-precision Pulsar Timing, MNRAS, 429, 2161
- Kramer, M. & Champion, D. J., 2013, The European Pulsar Timing Array and the Large European Array for Pulsars, CQGra, 30, 4009
- Lazio, T. J. W., 2013, The Square Kilometre Array Pulsar Timing Array, CQGra, 30, 4011
- Lentati, L., Alexander, P., & Hobson, M. P., et al., 2014, TEMPONEST: a Bayesian Approach to Pulsar Timing Analysis, MNRAS, 437, 3004
- Manchester, R., Hobbs, G. B., & Bailes, M., et al., 2013a, The Parkes Pulsar Timing Array Project, PASA, 30, 17
- Manchester, R., et al., 2013b, The International Pulsar Timing Array, CQGra, 30, 4010
- McLaughlin, M. A., 2013, The North American Nanohertz Observatory for Gravitational Waves, CQGra, 30, 4008
- Oslowski, S., van Straten, & Hobbs, G., et al., 2011, High Signal-to-noise Ratio Observations and the Ultimate Limits of Precision Pulsar Timing, MNRAS, 418, 1258
- Oslowski, S., van Straten, Demorest, P., W., & Bailes, M., 2013, Improving the Precision of Pulsar Timing Through Polarization Statistics, MNRAS, 430, 416
- Ravi, V., Wyithe, J. S. B., Shannon, R., & Hobbs, G., 2014a, MNRAS, 447, 2772
- Ravi, V., Wyithe, J. S. B., Shannon, R., Hobbs, G., & Manchester, R., 2014b, Binary Supermassive Black Hole Environments Diminish the Gravitational Wave Signal in the Pulsar Timing Band, MNRAS, 442, 56
- Sanidas, S. A., Battye, R. A., & Stappers, B. W., 2012, Constraints on Cosmic String Tension Imposed by the Limit on the Stochastic Gravitational Wave Background from the European Pulsar Timing Array, PhRvD, 85, 2003
- Sesana, A., Vecchio, A., & Colacino, C. N., 2008, The Stochastic Gravitational-wave Background from Massive Black Hole Binary Systems: Implications for Observations with Pulsar Timing Arrays, MNRAS, 390, 192
- Sesana A., Vecchio A., & Volonteri M., 2009, Gravitational Waves from Resolvable Massive Black Hole Binary Systems and Observations with Pulsar Timing Arrays, MNRAS, 394, 2255
- Shannon, R. M., Ravi, V., & Coles, W. A., et al., 2013, Gravitational-wave Limits from Pulsar Timing Constrain Supermassive Black Hole Evolution, Science, 342, 334
- Shannon, R. M., Oslowski, S., & Dai, S., et al., 2014, Limitations in Timing Precision due to Single-pulse Shape Variability in Millisecond Pulsars, MNRAS, 443, 1463
- van Haasteren, R., Levin Y., & Janssen, G., H., et al., 2011, Placing Limits on the Stochastic Gravitational-wave Background Using European Pulsar Timing Array Data, MNRAS, 414, 3117
- van Straten, W., 2013, High-fidelity Radio Astronomical Polarimetry Using a Millisecond Pulsar as a Polarized Reference Source, ApJS, 204, 13
- Verbiest, J. P. W., Bailes, M., & Coles, W. A., et al., 2009, Timing Stability of Millisecond Pulsars and Prospects for Gravitational-wave Detection, MNRAS, 400, 951
- Yan, W. M., Manchester, R. N., & van Straten, W., et al., 2011, Polarization Observations of 20 Millisecond Pulsars, MNRAS, 414, 2087
- Yardley, D. R. B., Coles, W. A., & Hobbs, G. B., et al., 2011, On Detection of the Stochastic Gravitational-wave Background Using the Parkes Pulsar Timing Array, MNRAS, 414, 1777
- Yardley, D. R. B., Hobbs, G. B., & Jenet, F. A., et al., 2010, The Sensitivity of the Parkes Pulsar Timing Array to Individual Sources of Gravitational Waves, MNRAS, 407, 669
- You, X. -P., Hobbs, G., Coles, W. A., Manchester, R. N., & Han, J. L., 2007, An Improved Solar Wind Electron Density Model for Pulsar Timing, ApJ, 671, 907
- You, X. -P., Coles, W. A., Hobbs, G., & Manchester, R. N., 2012, Measurement of the Electron Density and Magnetic Field of the Solar Wind Using Millisecond Pulsars, MNRAS, 422, 1160
- Zhu, X. -J. Hobbs, G., & Wen, L., et al., 2014, MNRAS, 444, 3709